

Instructions for taking MoviPrep and Senna Bowel Preparation



James Paget
University Hospitals
NHS Foundation Trust

Author: Valerie Dean, Radiographer

Why do I need to take the bowel preparation?

To ensure a thorough examination of your colon (large bowel) it must be emptied of waste material as your Radiographer or Radiologist needs clear views of the lining of your large bowel to enable detection of subtle abnormalities. The MoviPrep® bowel preparation is a strong laxative which will empty your bowel of waste material. It is important that you follow these instructions correctly. If you are not properly prepared the procedure may not be successful and may be cancelled.

Preparing for your Procedure:

An additional instruction sheet is given to people with diabetes to help them regulate their blood sugars during the bowel preparation and procedure. If you have diabetes and this has not been included please contact the Radiology Department. If you take any blood thinning medications (except Aspirin), please contact the Radiology Department Nurses on 01493 452099.

Seven days before the test

- Stop taking iron tablets.

Four days before the test

- Stop taking constipating medications, these may be painkillers such as codeine phosphate, tramadol, oramorph and fentanyl patches, bulking agents such as Fybogel®, Regulan ®or Proctofibe® or to treat diarrhoea, such as loperamide (Imodium®) or Lomotil®.
- If you are unable to stop taking your painkillers, please contact the Radiology department for advice.

Two days before the test

- **EAT FOODS LOW IN FIBRE.** These include grilled or steamed white fish, grilled chicken (no skin), potato (no skin), white pasta, white rice, egg, cheese, tofu, white bread, butter or margarine, seedless jam, shredless marmalade, marmite, honey, rich tea biscuits, chocolate, smooth yoghurts (no 'bits'), boiled sweets, clear jelly and ice cream.

AVOID HIGH FIBRE FOODS - Red meat, brown / wholemeal bread, wholemeal pasta, brown rice, fresh or dried fruit, vegetables, nuts, beans, lentils, chickpeas, cereals.

The day before the test

- Low fibre diet up until two hours before starting to take the MoviPrep®
- You are allowed clear fluids only until 2 hours before your appointment.

How to prepare Moviprep® solution

You have been given a pack containing four sachets of MoviPrep®, two X Sachet 'A' (light yellow powder) and two x Sachet 'B' (light brown powder) and eight Senna tablets (7.5mg)

To prepare the MoviPrep®:

- Empty the contents of one Sachet 'A' and one Sachet 'B' into a large jug.
- Add 1 litre (1¾ pts) of tap water.

- Stir until the powder is completely dissolved.
- The solution is now ready to drink.

When you start to drink your MoviPrep® solution you need to stay close to a toilet as you will experience frequent watery bowel movements. This is normal and indicates that the MoviPrep® solution is working. We recommend you use a barrier cream and wet wipes for your comfort.

You should start to have bowel movements one to two hours after starting to take MoviPrep®. You may experience some nausea (sickness) and stomach cramping. If you begin to vomit, please contact the Radiology department.

If you have not had any bowel movements by the day of your examination, contact the Radiology department for advice.

If you have a morning appointment

On the day before the test:

Before 8am Take all eight Senna tablets at one time.

After 3pm No further food can be eaten, you may continue to drink clear fluids.

Clear fluids allowed – water, clear soup (chicken, beef or vegetable strained to remove any bits), marmite, Bovril, clear fruit juice (no bits), soft drinks, fruit and herbal tea, tea without milk, coffee without milk.

Fluids not permitted – alcohol, milk or any drinks containing ‘bits’.

5pm Start drinking the first jug of MoviPrep®. Drink one glass every 15 minutes until you have drunk it all. Please drink another 600ml (1pt.) of clear fluids over this time to prevent dehydration.

8pm Repeat with the second set of sachets. You should have drunk the solution from all sachets within four – six hours. Please drink another 600ml (1pt.) of clear fluids over this time to prevent dehydration.

On the day of the test:

- No solid food
- You may drink clear fluids until two hours before your appointment.
-

If you have an afternoon appointment

On the day before the test:

Before 8am Take all eight Senna tablets at one time.

After 5pm No further food can be eaten, you may continue to drink clear fluids.

Clear fluids allowed – water, clear soup (chicken, beef or vegetable strained to remove any bits), marmite, Bovril, clear fruit juice (no bits), soft drinks, fruit and herbal tea, tea without milk, coffee without milk.

Fluids not permitted – alcohol, milk or any drinks containing ‘bits’.

7pm Start drinking the first jug of MoviPrep®. Drink one glass every 15 minutes until you have drunk it all. You should have drunk the solution within two hours. Please drink another 600ml (1pt.) of clear fluids over this time to prevent dehydration.

On the day of the test:

- No solid food
- **6am** Mix up and start drinking the second set of sachets. You should have drunk the solution within two hours. Please drink another 600ml (1pt.) of clear fluids over this time to prevent dehydration.
- You may drink clear fluids until two hours before your appointment.

Your Feedback

We want your visit to be as comfortable as possible - talk to the person in charge if you have any concerns. If the ward/department staff are unable to resolve it, then ask for our Patient Advice and Liaison (PALS) information. Please be assured that raising a concern will not impact on your care.

Before leaving please complete a Friends and Family Test feedback card.

Help us transform NHS services and to support patient choice.

OUR VALUES

Collaboration

We work positively with others to achieve shared aims

Accountability

We act with professionalism and integrity, delivering what we commit to, embedding learning when things do not go to plan

Respect

We are anti-discriminatory, treating people fairly and creating a sense of belonging and pride

Empowerment

We speak out when things don't feel right, we are innovative and make changes to support continuous improvement

Support

We are compassionate, listen attentively and are kind to ourselves and each other



 **IN TRAN**
communication for all

The hospital can arrange for an interpreter or person to sign to assist you in communicating effectively with staff during your stay. Please let us know.

For a large print version of this leaflet, contact PALS 01493 453240